

SEASONAL GROWTH RATES OF FISHES IN RELATION TO CONDITIONS OF LAKE STRATIFICATION

Glen E. Gebhart and Robert C. Summerfelt*

Oklahoma Cooperative Fishery Research Unit[†], Oklahoma State University, Stillwater, Oklahoma

The effect of lake stratification and hypolimnetic anoxia on growth of gizzard shad (*Dorosoma cepedianum*), white crappie (*Pomoxis annularis*), and freshwater drum (*Aplodinotus grunniens*) was studied in Arbuckle Lake, Oklahoma. The percentage of total available fish habitat of Arbuckle Lake was decreased by anoxic conditions to 72.9% in 1973, 82.7% in 1974, and 72.7% in 1975. In 6 of 8 species-age group comparisons, the growth rate was highest in 1974 when more usable fish habitat was available than in other years. In 18 species-age group comparisons of fish growth rates in the summer stratified period with growth of the same age fish in the destratified overwinter period, the growth was faster in the destratified interval 12 times and faster in the stratified interval 6 times. It appears that lake stratification and the resultant loss of fish habitat retarded the growth of the fish.

INTRODUCTION

The objective of this paper is to examine the effect of lake stratification on fish growth. Most lakes and reservoirs in the United States are stratified in the summer, and dissolved oxygen (DO) in the hypolimnion is often depleted. Anoxia restricts fish from continuous occupancy of the hypolimnion of stratified lakes and reservoirs (1, 2). During middle and late summer, stratification commonly causes compression of even warm-water fishes into the stratum of water between the anoxic hypolimnion and the warm epilimnion where surface temperatures often exceed their thermal preferences. The compression effect of stratification may be lethal to cold-water fishes in lakes lacking cold, well-oxygenated water (3).

In Red Haw Lake, Iowa, Mayhew (4) noted that at the time of severe stratification and hypolimnetic stagnation most scales of bluegills (*Lepomis macrochirus*) developed false annuli, and their growth was greatly retarded in relation to the prestratification rate. Mayhew suggested that stratification is a primary controlling factor in growth of bluegills. Seasonal periodicity of growth has been reported which shows a reduced rate in midsummer. This coincides with the onset of summer stratification and severe hypolimnetic oxygen depletion. Beckman (5) found that 85% of the expected growth of bluegills in a Michigan lake was completed between the end of April and mid-July. Sprugel (6) found bluegill growth in Iowa was most rapid in June and little growth evident after July. Anderson (7) also found bluegill growth to be most rapid in June with most growth occurring in May, June and July and the total growing season extending from 1 May to 31 October. Winter temperatures are generally too low for optimum growth of most warm-water fishes, but even normal summer surface temperatures can exceed the optimum range for warm-water fishes as ascertained from laboratory and field studies of the preferred temperatures of fish (8).

In spite of an abundant literature on growth of fish, Mayhew's report remains the only direct examination of fish growth in relation to conditions of stratification. Since stratification is a well known limnological phenomenon having substantial ecological importance (9), it is surprising to find the lack of direct investigations relating stratification to fish growth.

The inimical water quality conditions occurring during stratification (high ammonia, hydrogen sulfide, and carbon dioxide and low DO) and their concomitant effects on production of benthic invertebrates are likely to retard the growth rates of fishes. Laboratory experiments in which conditions in stratified lakes are simulated show diminished fish growth rates at low DO levels and high tempera-

*Present address: Department of Animal Ecology, Iowa State University, Ames, Iowa 50011.

[†]Cooperators are the Oklahoma Department of Wildlife Conservation, the Oklahoma State University Environmental Institute, and the U.S. Fish and Wildlife Service.

tures. Stewart et al. (10) reported that growth rates of juvenile largemouth bass (*Micropterus salmoides*) were markedly reduced at DO levels below air saturation. The growth of bass subjected to fluctuating oxygen concentrations was also markedly impaired. Andrews et al. (11) found a similar decreased growth of channel catfish (*Ictalurus punctuatus*) held at reduced DO levels. Andrews and Stickney (12) observed a reduction in the growth rate of channel catfish when the temperature was increased from 30 C to 34 C.

Arbuckle Lake, Oklahoma, is typically stratified from mid-May through September with anoxic conditions occurring below 5 to 8 m (13). In each of the three years of our study, 1973-75, Arbuckle Lake was thermally stratified by mid-May and DO depletion followed the onset of thermal stratification (2). The lake remained stratified until September or October depending on the weather and on the artificial mixing which was attempted in these years. During the summer of 1973, the Bureau of Reclamation attempted destratification by using a compressed air gun similar to one described by Bryan (14). During the summers of 1974 and 1975 an attempt was made by J. E. Garton of Oklahoma State University to destratify the reservoir using an axial flow pump, larger than but similar to one described by Quintero and Garton (15).

The present study describes seasonal growth rates of gizzard shad (*Dorosoma cepedianum*), white crappie (*Pomoxis annularis*), and freshwater drum (*Aplodinotus grunniens*) in Arbuckle Lake from collections made in 1973-75.

METHODS

In 1973, fish collections were made for 15 continuous weeks from 14 May to 24 August and an additional collection was obtained 18-21 October. In 1974, collections began on 10-14 March 1974 with the weekly collection scheduled beginning 20 May and continuing through 23 August. A final 1974 collection was made 9-13 September to monitor growth and distribution of fish after the lake had destratified. In 1975, weekly collections began on 12 May and continued through 22 August; and a collection was made 22-26 September after the lake had destratified. The collection of fish at the beginning and end of the stratified period allowed for the measurement of growth during both the stratified and destratified periods.

Fish were sampled with experimental gill nets, 3.05 m deep by 45.73 m long, divided into six 7.62-m panels with mesh sizes (bar measure) of 1.27, 2.54, 3.81, 5.08, 6.35, and 7.62 cm. Nets were set in a horizontal position at depths of 0-5, 5-10, 10-15, and 15-20 m. The vertical distribution of fish, described from these net catches, showed the expected movement of fish out of the hypolimnion when it became anoxic (2). Weights from fish of the same age, collected at four sites, were pooled to obtain a mean weight for each period. Fish were weighed and measured, and scales were taken to determine the age of the fish by annular counts. Temperature and DO measurements were taken weekly at 1-m intervals at all sites with a DO probe and thermistor.

Instantaneous growth rates (G) were calculated for the stratified summer and the destratified overwintering periods for all age groups of the three species that were adequately represented in the sample. Values of G were calculated by the equation of Chapman (16). The growth rates and mean weight gains (g) were calculated from samples collected at the start and end of the stratified period.

RESULTS

In 7 of 8 destratified vs. stratified comparisons of G and g for gizzard shad, larger values were obtained in the destratified period (Table 1). Only in 1973, for age IV gizzard shad, were the G and g values larger in the stratified period. The results of t-tests showed significantly higher growth rates ($P < .05$) and mean weight gains ($P < .01$) of all age groups of gizzard shad combined in the destratified period than in the summer stratified period.

The mean weight gains for age group I white crappie were higher in the destratified period than in the stratified period in 2 of 3 yearly comparisons (Table 2). The G values for age I white crappie were higher in the stratified period in 2 of 3 years. Age II white crappie had larger absolute growth in the stratified interval in 2 of the 3 years, but higher growth rates in the destratified interval in 2 of 3 years.

TABLE 1. Comparison of growth rates (G) and mean weight gain for gizzard shad for summer (stratified) and winter (destratified) intervals, 1973-76.

Median collection date	Interval	Days	Age group III			Age group IV			Age group V		
			Growth rate	Weight gain (g)	Sample size	Growth rate	Weight gain (g)	Sample size	Growth rate	Weight gain (g)	Sample size
5-23-73	summer	149.0	-0.17	-2.0	237	1.33	29.2	88	-0.13	-3.8	21
10-19-73	winter	221.5	1.58	31.1	92	0.53	20.5	46	0.33	14.5	26
5-29-74	summer	105.5	0.98	8.7	87	1.40	16.8	10	-0.79	-14.7	25
9-11-74	winter	252.0	1.20	31.4	47	1.69	65.1	114	—	—	—
5-21-75	summer	128.0	0.71	7.2	48	1.13	18.8	39	-1.72	-37.0	120
9-24-75	winter	236.5	2.96	83.8	46	1.24	47.6	63	1.61	69.9	78

TABLE 2. Comparison of growth rates (G) and mean weight gain for white crappie for summer (stratified) and winter (destratified) intervals, 1973-76.

Median collection date	Interval	Days	Age group I			Age group II			Age group III		
			Growth rate	Weight gain (g)	Sample size	Growth rate	Weight gain (g)	Sample size	Growth rate	Weight gain (g)	Sample size
5-23-73	summer	149.0	6.30	21.7	6	1.42	30.2	50	0.08	2.9	26
10-19-73	winter	221.5	0.90 ^a	19.7	19	2.06	102.0	5	—	—	—
5-29-74	summer	105.5	8.25	42.8	102	6.37	142.3	16	2.39 ^b	74.2	6
9-11-74	winter	252.0	4.39	149.0	78	0.64	51.7	11	—	—	—
5-21-75	summer	128.0	0.12	0.8	167	-0.90	-24.4	56	1.54 ^c	34.3	5
9-24-75	winter	236.5	1.35	19.4	255	-0.82	-35.0	45	—	—	—

^aInterval was 241.5 days in length.

^bInterval was 85.5 days in length.

^cInterval was 62.0 days in length.

2 mg/l DO (2). The percentage of the total volume of lake water with less than 2 mg/l DO was as high as 48% in 1973, 39% in 1974, and 59% in 1975. The average volume of stored water containing more than 2 mg/l DO during the 163-day interval from 11 May to 20 October (during a typical year this interval should encompass most of the summer stratification period) was 72.9% in 1973, 82.7% in 1974, and 72.7% in 1975. There was little difference between 1973 and 1975, but the percentage in both these years differed substantially from that in 1974 (2).

This variation in available habitat was reflected in yearly variations in the growth rate of fish during the stratified period. The fastest growth of age group III and IV gizzard shad and the second fastest growth for age V shad were observed in 1974 (Table 1). All three age groups of white crappies experienced greater growth in 1974 than in the other years (Table 2). The growth of age group II freshwater drum in 1974 was second to growth in 1975 but the maximum growth rate of age III drums was in 1974 (Table 3).

It appears that the seasonal cycle of stratification reduces fish growth by reducing the amount of fish habitat in a lake. The growth of fishes appears to slow down when lakes develop an anoxic hypolimnion due to stratification. Fish forced out of the cooler waters of the hypolimnion must remain in the transitional area of the thermocline if epilimnetic temperatures are thermally oppressive; or if they tolerate the higher temperatures of the epilimnion, food consumption must be increased to maintain a higher basal metabolism. It seems that the food supply is generally inadequate to provide for their basal metabolism needs, since many age groups lose weight during the summer.

REFERENCES

1. J. S. DENDY, J. Tenn. Acad. Sci. 20: 114-135 (1945).
2. G. E. GEBHART and R. C. SUMMERFELT, J. Envir. Eng. Div. ASCE 102 (EE6): 1215-1228 (1976).
3. P. J. COLBY and L. T. BROOKE, Limnol. Oceanogr. 14: 958-960 (1969).
4. J. MAYHEW, Biol. Sec. Rep. Iowa Conserv. Comm., 1963.
5. W. C. BECKMAN, Trans. Am. Fish. Soc. 72: 72-78 (1943).
6. G. SPRUGEL, JR., Trans. Am. Fish. Soc. 83: 58-75 (1954).
7. R. D. ANDERSON, *The Influence of Season and Temperature on Growth of the Bluegill, Lepomis Macrochirus Rafinesque*, Ph.D. Thesis, Univ. Michigan, Ann Arbor, 1959.
8. R. G. FERGUSON, J. Fish. Res. Board Can. 15: 607-624 (1958).
9. D. TOETZ, R. C. SUMMERFELT, and J. WILHM, *Biological Effects of Artificial Destratification in Lakes and Reservoirs — Analysis and Bibliography*, Rep. REC-ERC-72-33, U.S. Bur. Reclamation, Denver, 1972.
10. N. E. STEWART, D. L. SHUMWAY, and P. DOUDOROFF, J. Fish. Res. Board Can. 24: 475-494 (1967).
11. J. W. ANDREWS, T. MURAI, and G. GIBBONS, Trans. Am. Fish. Soc. 102: 835-838 (1973).
12. J. W. ANDREWS and R. R. STICKNEY, Trans. Am. Fish. Soc. 101: 94-99 (1972).
13. W. R. DUFFER and C. G. HARLIN, JR., *Changes in Water Quality Resulting from Impoundment*, U.S. Environ. Prot. Agency Water Pollut. Control Res. Ser. Proj. No. 16080 GGH, 1971.
14. J. G. BRYAN, *Symposium on Streamflow Regulation for Quality Control*, U.S. Public Health Serv. Publ. No. 999-WP-30, Cincinnati, 1965, pp. 317-334.
15. J. E. QUINTERO and J. E. GARTON, Trans. Am. Soc. Agric. Eng. 16: 973-978 (1973).
16. D. W. CHAPMAN, in W. E. RICKER (ed.), *Methods for Assessment of Fish Production in Fresh Waters*, Int. Biol. Program Handbook No. 3, Blackwell Scientific Publications, Oxford, 1971, pp. 199-214.
17. F. W. TESCH, in: W. E. RICKER (ed.), *Methods for Assessment of Fish Production in Fresh Waters*, Int. Biol. Program Handbook No. 3, Blackwell Scientific Publications, Oxford, 1971, pp. 93-130.